ELSEVIER

Aquaculture 216 (2003) 391–393

Aquaculture

www.elsevier.com/locate/aqua-online

Contents of *Aquaculture*, Volume 216

VOL. 216 NOS. 1–4 10 February 2003

Diseases, Parasites and Predators

Primary cultures of hemocytes from *Mytilus galloprovincialis* Lmk.: expression of IL-2Rα subunit
A. Cao (Lugo, Spain), L. Mercado (Chile), J.I. Ramos-Martinez and R. Barcia (Lugo, Spain) 1

Qualitative and quantitative studies on the relative virus load of tails and heads of shrimp acutely infected with WSSV
S.V. Durand (Tucson, AZ, USA; Montpellier, France), R.M. Redman, L.L. Mohney, K. Tang-Nelson (Tucson, AZ, USA), J.R. Bonami (Montpellier, France) and D.V. Lightner (Tucson, AZ, USA) 9

Induced resistance to white spot syndrome virus infection in *Penaeus stylirostris* through pre-infection with infectious hypodermal and hematopoietic necrosis virus—a preliminary study
K.F.J. Tang, S.V. Durand, B.L. White, R.M. Redman, L.L. Mohney and D.V. Lightner (Tucson, AZ, USA) 19

Effect of fasting on feed intake, growth and mortality of chinook salmon, *Oncorhynchus tshawytscha*, during an induced *Aeromonas salmonicida* epizootic
J. Pirhonen, C.B. Schreck (Corvallis, OR, USA), P.W. Reno (Newport, OR, USA) and H. Ögüt (Trabzon, Turkey) 31

Genetics and Breeding

Effects of heat-shock on cell division and microtubule organization in zygotes of the shrimp *Penaeus indicus* (Crustacea, Decapoda) observed with confocal microscopy
M. Morelli (Nouméa, New Caledonia), Aquacop (Taravao, French Polynesia) 39

Production of a high percentage of male offspring with a natural androgen, 11β-hydroxyandrostenedione (11βOHA4), in Florida red tilapia
D. Desprez (Montpellier, France; Belgium), E. Géraz (Montpellier, France), M.C. Hoareau (France), C. Mélard (Belgium), P. Bosc (France) and J.F. Baroiller (Montpellier, Rennes, France) 55

A combined salt and confinement stress enhances mortality in rainbow trout (*Oncorhynchus mykiss*) selected for high stress responsiveness
S.-E. Fevolden (Tromsö, Norway), K.H. Roed (Oslo, Norway) and K. Fjalestad (Norway) 67

Husbandry and Management

Identification and growth rates comparison of divergent migratory contingents of Japanese eel (*Anguilla japonica*)
W.-N. Tzeng, Y. Iizuka, J.-C. Shiao (Taipei, Taiwan), Y. Yamada and H.P. Oka (Japan) 77

A beneficial coculture: charm abalone *Haliotis discus hannai* and sea cucumber *Stichopus japonicus*
K.H. Kang, J.Y. Kwon and Y.M. Kim (Yosu, South Korea) 87

Transport of silver catfish (*Rhamdia quelen*) fingerlings at different times, load densities, and temperatures
J.I. Golombieski, L.V.F. Silva, B. Baldisserotto and J.H.S. da Silva (Santa Maria, Brazil) 95

Do stocking density and feed reward level affect growth and feeding of self-fed juvenile European sea bass?
M. Paspatis (Heraclion, Greece), T. Boujard (France), D. Maragoudaki (Heraclion, Greece), G. Blanchard (Namur, Belgium) and M. Kentouri (Heraclion, Greece) 103
Variability in mussel settlement on suspended ropes placed at Ahipara Bay, Northland, New Zealand
A.C. Alfaro and A.G. Jeffs (Auckland, New Zealand) 115
Growth and survival of the tehuelche scallop *Aequipecten tehuelchus* in culture
M.A. Narvarte (Argentina) 127
Bioencapsulation in *Artemia*: II. Influences of the particle concentration in the enrichment process
R. Gelabert F. (Ciudad del Carmen, Mexico) 143
KCl induced paralysis facilitates detachment of hatchery reared juvenile green sea urchins, *Strongylocentrotus droebachiensis*
N.T. Hagen (Bodø, Norway) 155
Food intake of *Macrobrachium rosenbergii* during larval development
H.P. de Barros and W.C. Valenti (São Paulo, Brazil) 165

Nutrition

The resistance to physical stresses by *Penaeus monodon* juveniles fed diets supplemented with astaxanthin
Y.-H. Chien, C.-H. Pan (Keelung, Taiwan) and B. Hunter (Bangkok, Thailand) 177
Use of the bacteria *Streptococcus faecium* and *Lactobacillus acidophilus*, and the yeast *Saccharomyces cerevisiae* as growth promoters in Nile tilapia (*Oreochromis niloticus*)
M. Lara-Flores, M.A. Olvera-Novoa, B.E. Guzmán-Méndez and W. López-Madrid (Mérida, Mexico) 193
Optimisation of T-ISO biomass production rich in essential fatty acids: I. Effect of different light regimes on growth and biomass production
I. Tzovenis, N. De Pauw and P. Sorgeloos (Ghent, Belgium) 203
Optimisation of T-ISO biomass production rich in essential fatty acids: II. Effect of different light regimes on the production of fatty acids
I. Tzovenis, N. De Pauw and P. Sorgeloos (Ghent, Belgium) 223
Optimum daily ration for Eurasian perch *Perca fluviatilis* L. reared at its optimum growing temperature
E.D. Fiogbé (Cotonou, Benin) and P. Kestemont (Namur, Belgium) 243
Apparent digestibility of selected feedstuffs by mud crab, *Scylla serrata*
M.R. Catacutan, P.S. Eusebio (Tigbauan, Philippines) and S.-i. Teshima (Kagoshima City, Japan) . . 253
Determination of fat content in fish feed by supercritical fluid extraction and subsequent lipid classification of extract by thin layer chromatography-flame ionization detection
R.B. Johnson and H.J. Barnett (Seattle, WA, USA) 263
Potato protein concentrate with low content of solanidine glycoalkaloids in diets for Atlantic salmon (*Salmo salar*)
S. Refstie (Sunndalsøra, Norway) and H.A.J. Tiekstra (Veendam, The Netherlands) 283
Lipid and fatty acid composition of ovaries from wild fish and ovaries and eggs from captive fish of white sea bream (*Diplodus sargus*)
J.R. Cejas, E. Almansa, J.E. Villamandos, P. Badia, A. Bolaños and A. Lorenzo (Santa Cruz de Tenerife, Spain) 299

Physiology and Endocrinology

Gonadal differentiation and effects of temperature on sex determination in southern flounder (*Paralichthys lethostigma*)
J.A. Luckenbach, J. Godwin, H.V. Daniels and R.J. Borski (Raleigh, NC, USA) 315
Expression of common carp growth hormone in the yeast *Pichia pastoris* and growth stimulation of juvenile tilapia (*Oreochromis niloticus*)
Y. Li, J. Bai, Q. Jian, X. Ye, H. Lao, X. Li, J. Luo and X. Liang (Guangzhou, PR China) 329
Influences of light intensity and photoperiod on moulting and growth of *Penaeus merguiensis* cultured under laboratory conditions
T. Hoang, M. Barchiesis, S.Y. Lee, C.P. Keenan and G.E. Marsden (Grafton, Australia) 343

Differences in the biochemical content of buoyant and non-buoyant eggs of the Japanese eel, *Anguilla japonica*
M. Seoka, S. Yamada, Y. Iwata, T. Yanagisawa, T. Nakagawa and H. Kumai (Wakayama, Japan) . . 355
Brain activity of Atlantic salmon (*Salmo salar*) following electrical stunning using various field strengths and pulse durations
D.H.F. Robb (Bristol, UK) and B. Roth (Bergen, Norway) 363
Induction of sperm activation in open and closed thelycum penaeoid shrimps
J. Alfaro, N. Muñoz (Puntarenas, Costa Rica), M. Vargas (San José, Costa Rica) and J. Komen (Wageningen, Netherlands) 371

Book review
The biology of decapod crustacean larvae (Crustacean Issues 14)
M.B. New (Marlow, UK) 383

Erratum
Erratum to "Genetic variation for growth rate, feed conversion efficiency, and disease resistance exists within a farmed population of rainbow trout" [Aquaculture 209 (2002) 59–76]
M. Henryon, A. Jokumsen, P. Berg, I. Lund, P.B. Pedersen, N.J. Olesen and W.J. Slierendrecht (Aarhus, Denmark) 389

Contents of *Aquaculture*, Volume 216 391